

Certificate of Analysis

Page 1 of 3

| | | | | |
|-----------------|---|--------------------------|--------------------------------|-------|
| Client: | Cosana New Zealand Limited | Lab No: | 3892539 | HGPV1 |
| Contact: | Cherie Kloet C/- Cosana New Zealand Limited PO Box 3330 Taupo 3351 | Date Received: | 19-May-2025 | |
| | | Date Reported: | 23-May-2025 | |
| | | Quote No: | 130194 | |
| | | Order No: | SORD25117 | |
| | | Client Reference: | CSNZ14867 - RARNZ - Batch Test | |
| | | Submitted By: | Cherie Kloet | |

| Sample Type: Honey | | | | |
|--|---------------------|-------------------------|--|--|
| Sample Name: | | CSNZ14867 | | |
| Lab Number: | | 3892539.1 | | |
| MPI Manuka Classification | | | | |
| MPI Manuka Honey Classification | | Monofloral Manuka Honey | | |
| 3-Phenyllactic acid (3-PA) | mg/kg | 1,090 | | |
| 2'-Methoxyacetophenone (2'-MAP) | mg/kg | 19.3 | | |
| 2-Methoxybenzoic acid (2-MBA) | mg/kg | 17.4 | | |
| 4-Hydroxyphenyllactic acid (4-HPA) | mg/kg | 11.0 | | |
| Manuka DNA | Cq | 34.22 | | |
| Manuka Honey Analysis | | | | |
| Dihydroxyacetone (DHA) | mg/kg | 1,174 | | |
| 5-Hydroxymethylfurfural (HMF) | mg/kg | 37.4 | | |
| Methylglyoxal (MGO) | mg/kg | 871 | | |
| Non Peroxide Activity (NPA)* | % Phenol Equivalent | 20.6 | | |
| Leptosperin | mg/kg | 370 | | |
| Tutin Analysis | | | | |
| Tutin Result Evaluation | Pass/Fail | PASS | | |
| Tutin | mg/kg | < 0.010 | | |
| MRL as per Tutin in Honey Food Standard 2016 | mg/kg | 0.70 | | |
| Microbiological Analysis | | | | |
| Aerobic Count 35°C | cfu / g | < 10 | | |
| Yeasts & Moulds | cfu / g | < 10 | | |
| Total Coliforms | cfu / g | < 10 | | |
| Staphylococcus aureus | cfu / g | < 10 | | |
| Physical Analysis | | | | |
| Moisture | g/100g as rcvd | 17.3 | | |
| Glyphosate Analysis | | | | |
| AMPA | mg/kg | < 0.010 | | |
| Glufosinate | mg/kg | < 0.010 | | |
| Glyphosate | mg/kg | < 0.010 | | |

Analyst's Comments

Sample 1 Comment:

MPI Classification Comment:

The results presented on the Certificate of Analysis have been rounded to an appropriate number of significant figures, based on the Uncertainty of Measurement of the methods performed. The 'MPI Manuka Honey Classification' has been determined using unrounded values. In cases where one or more values were close to the critical levels (as defined by MPI), there may be a seeming inconsistency between the classification and the rounded values reported.

Summary of Methods

The following table(s) gives a brief description of the methods used to conduct the analyses for this job. The detection limits given below are those attainable in a relatively simple matrix. Detection limits may be higher for individual samples should insufficient sample be available, or if the matrix requires that dilutions be performed during analysis. A detection limit range indicates the lowest and highest detection limits in the associated suite of analytes. A full listing of compounds and detection limits are available from the laboratory upon request. Unless otherwise indicated, analyses were performed at Hill Labs, 28 Duke Street, Frankton, Hamilton 3204.

Sample Type: Honey

| Test | Method Description | Default Detection Limit | Sample No |
|---------------------------------------|---|-------------------------|-----------|
| Individual Tests | | | |
| Moisture | Refractometer. Harmonised Methods of the International Honey Commission (2009). Method 1 - Determination of Moisture, Refractometric Method. (modified). | 13.0 g/100g as rcvd | 1 |
| 3-in-1 Honey method | Aqueous extraction, derivatisation. Analysis by uHPLC / UV-Vis (dihydroxyacetone, 5-hydroxymethylfurfural, methylglyoxal). In-house. | 1.0 - 10 mg/kg | 1 |
| Leptosperin | Aqueous extraction, dilution, analysis by LC-MS/MS. | 15 mg/kg | 1 |
| Non Peroxide Activity (NPA)* | NPA is calculated from methylglyoxal using an industry accepted correlation curve based on published data ^{1,2} for NPA and the primary active ingredient, methylglyoxal. ¹ Isolation by HPLC and characterisation of the bioactive fraction of New Zealand manuka (<i>Leptospermum scoparium</i>) honey. C. J. Adams, et al. Carbohydrate Research 343 (2008) 651-659. ² Corrigendum to "Isolation by HPLC and characterization of the bioactive fraction of New Zealand manuka (<i>Leptospermum scoparium</i>) honey" [Carbohydr. Res. 343 (2008) 651]. C. J. Adams, et al. Carbohydrate Research 344 (2009) 2609. | 1.0 % Phenol Equivalent | 1 |
| Tutin Analysis in Honey | Solvent extraction, dilution. Analysis by LC-MS/MS. Results are representative of the liquid honey, not the sample as a whole. <u>Tutin Result Evaluation (PASS/FAIL)</u> The PASS/FAIL result is based on comparison of the tutin result with the "Food Standard: Tutin in Honey (2016)". A result that falls at or BELOW the maximum permitted tutin level will give a PASS result. A result that falls ABOVE the maximum permitted tutin level will give a FAIL result. <u>Individual Sample Testing Recommended?</u> Where a tutin result for a composited sample is above the maximum permitted level, it is recommended that the individual samples are retested. Please contact the laboratory to arrange for individual sample retesting. RLP Official Test 8.42. | 0.010 mg/kg | 1 |
| Aerobic Count 35°C | Automated MPN count on TEMPO AC, Incubated at 35°C for 22-28 hours. bioMérieux, TEMPO. | 10 cfu / g | 1 |
| Total Coliforms | Automated MPN count on TEMPO TC, incubated at 35°C for 24-27 hours. bioMérieux, TEMPO. | 10 cfu / g | 1 |
| Staphylococcus aureus | Automated MPN count on TEMPO STA, Incubated at 35°C for 24-27 hours. bioMérieux, TEMPO. | 10 cfu / g | 1 |
| Yeasts & Moulds | Automated MPN count on TEMPO YM, Incubated at 25°C for 72-76 hours. bioMérieux, TEMPO. | 10 cfu / g | 1 |
| Glyphosate LC-MS/MS Analysis | Aqueous extraction, Analysis by LC-MS/MS. In-house. RLP Official Test 8.47.1. | 0.010 mg/kg | 1 |
| MPI 5 Attributes Tests | | | |
| MPI Manuka Honey Classification | Evaluation of results against Ministry of Primary Industries (MPI) criteria for classification of monofloral and multifloral Manuka honey. General Export Requirements for Bee Products - 27 October 2021. | - | 1 |
| Manuka Honey Chemistry Profile | | | |
| 3-Phenyllactic acid (3-PA) | Aqueous solvent extraction, dilution. LC-MS/MS analysis. MPI Technical Paper 2017/30 (modified) RLP Official Test 10.05. | 5 mg/kg | 1 |
| 2'-Methoxyacetophenone (2'-MAP) | Aqueous solvent extraction, dilution. LC-MS/MS analysis. MPI Technical Paper 2017/30 (modified) RLP Official Test 10.05. | 0.50 mg/kg | 1 |

| Sample Type: Honey | | | |
|------------------------------------|---|-------------------------|-----------|
| Test | Method Description | Default Detection Limit | Sample No |
| 2-Methoxybenzoic acid (2-MBA) | Aqueous solvent extraction, dilution. LC-MS/MS analysis. MPI Technical Paper 2017/30 (modified) RLP Official Test 10.05. | 0.50 mg/kg | 1 |
| 4-Hydroxyphenyllactic acid (4-HPA) | Aqueous solvent extraction, dilution. LC-MS/MS analysis. MPI Technical Paper 2017/30 (modified) RLP Official Test 10.05. | 0.50 mg/kg | 1 |
| Manuka Honey PCR Profile | | | |
| Manuka DNA | Quantification of Manuka (<i>Leptospermum scoparium</i>) DNA by real time PCR. MPI Technical - Paper No: 2017/31 (modified). RLP Official Test 10.04. | > 36 Cq | 1 |

These samples were collected by yourselves (or your agent) and analysed as received at the laboratory.

Testing was completed between 20-May-2025 and 23-May-2025. For completion dates of individual analyses please contact the laboratory.

Samples are held at the laboratory after reporting for a length of time based on the stability of the samples and analytes being tested (considering any preservation used), and the storage space available. Once the storage period is completed, the samples are discarded unless otherwise agreed with the customer. Extended storage times may incur additional charges.

This certificate of analysis must not be reproduced, except in full, without the written consent of the signatory.

Shaun Clay BSc
Senior Technologist - Food and Bioanalytical